

Light-induced anomalous Hall effect in graphene

James W. McIver

Max Planck Institute for the Structure and Dynamics of Matter, Hamburg, Germany

Many non-equilibrium phenomena have been discovered or predicted in quantum solids driven by femtosecond pulses of light. Examples include photo-induced superconductivity [1] and Floquet-engineered topological phases [2]. These effects are expected to lead to measurable changes in electrical transport. However, the timescales involved far exceed those accessible using conventional fast electronics.

In this talk, I will present results on the transport properties of monolayer graphene illuminated by a femtosecond pulse of circularly polarized light [3]. This was achieved using an ultrafast device architecture based on laser-triggered photoconductive switches. We observed a light-induced Hall effect in the absence of an applied magnetic field. The dependence of the effect on a gate potential used to tune the Fermi level revealed multiple features that reflect a Floquet-engineered topological band structure, similar to the band structure originally proposed by Haldane [4]. This includes a ~ 60 meV wide conductance plateau centered at the Dirac point, where a gap of equal magnitude is predicted to open based on Floquet theory. We find that when the Fermi level lies within this plateau, the non-equilibrium anomalous Hall conductance saturates around $\sim 1.8 \pm 0.4 e^2/h$.




Fig. 1: Schematic of the device architecture used to detect ultrafast light-induced anomalous Hall currents in graphene.

References

- [1] D. Fausti et al. *Science* **331**, 189-191 (2011)
- [2] T. Oka & H. Aoki. *Phys. Rev. B* **79**, 081406 (2009)
- [3] J.W. McIver et al. *Nature Physics* (forthcoming); arXiv:1811.03522 (2019)
- [4] F.D.M. Haldane, *Phys. Rev. Lett.* **61**, 2015-2018 (1988)